References:

1. Larry JC and Herbert AL (1987), Suppositories. In: Lachman L, Liberman HA and Kanig JC. (eds.) *The Theory and Practice of Industrial Pharmacy.* 3rd ed. Varghese Publishing House, Mumbai 564-587.
2. Lawrence HB. Remington (1995), *The Science and Practice of Pharmacy*. 19th ed. Mack Publishing House, New York 1577.
3. Rawlins, E.A.(2002) Bentley's Text Book of Pharmaceutics 8th ed. Delhi, All India Traveller Book Seller, pp. 347-50
4. Rytting, J.H., Fix, J.A. (2002) Encyclopedia of Pharmaceutical Technology 2nd ed,New York, Marcel Dekker Inc, pp. 932-43
5. Hermann, T.W. (1995) Recent research on bioavailability of drugs from suppositories. Int. J. Pharm, 123., 1-11.
6. Sweetman S.C. (2005) Martindale: The Complete Drug Reference (34th ed., p.11). London, Pharmaceutical Press 11 & 865.
7. Akala EO, Adedoyn A and Ogunbona FA(1991). Suppository formulation of amodiaquine *in vitro* release characteristics. Drug Develop Ind Pharm, 17(2)., 303-07.
8. Allen LV (1997). Suppositories as drug delivery system Part-2. Journal of Pharmaceutical Care in Pain and Symptom Control.5(2)., 17-26.
9. Asikoglu M, Ertan G, Cosar G(1995). The release of isoconazole nitrate from different suppository bases: *In vitro* dissolution, physicochemical and microbiological studies. J. Pharm. Pharmacol, 47(9)., 713-6.
10. Basavaraj BV, Sindhu Abraham, Devaswaran R, Bharat S, Madhavan V(2007). Mucoadhesive effect of carbopol on drug release rate from double phased suppositories. Asian J Pharm, 1(1)., 93-5.
11. Brummer JM, Schoenmakers EAJM, Kumerink GJ, Heidendal GAK, Sander DGM, Stockbrugger RW(1997). The effect of single rectal dose of cisapride on delayed gastric emptying. Aliment Pharmacol Ther, 11(4)., 781-85.
12. Dash AK and Cudworth GC(2001). Evaluation of an acetic acid ester of monoglyceride as a suppository base with unique properties. AAPS Pharmascitech, 2(3)., Article 13. Available from url: http:// www.AAPS pharmascitech.com.

13. El Assasy, Foda AH, Badawi SS, Abdel Rahim RT(1995), Release characteristics and bioavailability of pirprofen from suppository bases. Egyptian J Pharm Sci, 36(1-6)., 15-25.
14. Elbary AA, El Nabarawi MA, Mohamed MI(1998). Effect of chemical structure on the release of certain propionic acid derivatives from their dosage forms. Drug Develop Ind Pharm, 24(5)., 439-451.
15. Taha EI, Zaghoul AA, SAmy AM, Al-Saidan S, Kassem AA, Khan MA(2004), Bioavailability assessment of salbutamol sulfate in human volunteers, Int.J.Pharm, 279., 3-7
16. Faruk Ahmed, Ashraful Islam Sm, Qadir MA, Saleem Reza MD.(2004). Soluble drug theophylline sodium glycinate from different suppository bases, Dhaka Univ.J.Pharm.Sci, 3(1-2).,
17. Hender T, Hender J, Gellin F, Haung ML, Vande PS, Woestenborghs R et al(1990). Comparative bioavailability of cisapride suppository and tablet formulation in healthy volunteers. E J Clin Pharmacol, 38(6)., 629-31.
18. Hudson Kristofer C, Asbill C Scott(2007), Webster Andrew A. Isoniazid Release from suppositories compounded with selected bases. Veternary Compounding., 15
19. Iwata M, Takayama K, Takahashi Y, Obtata Y, Shirotaka S(1998). Release of diclofenac sodium from suppositories consisting of witepsols as a base. Japanese J Hosp Pharm, 24(4)., 3587-62.
20. Jawahar N, Jayaprakash S, Maria Gerald Rajan NS, Nagarajan M, Dhachina Murthi D, Jibie S et .al (2005). Design and evaluation of sustained release suppositories of nimesulide. Indian J Pharm Sci, 67(5)., 558-61.
21. Kamlinder KS, Deshpande SG and Baichwal MR (1994). Studies on suppository bases: Design and Evaluation of Sodium CMC and Agar bases. Indian Drugs, 31., 149-54.
22. Khamaruzzaman MA, Ashraful islam SM, JAkir AC, Selim Reza MD(2005). Effect of Viscosity Imparting Agents on In-Vitro Drug Release from PEG Based Suppositories, Dhaka.Univ.J.Pharm, 4(1).,
23. Kim JY, Ku YS (2000). Enhanced absorption of indomethacin after oral or rectal administration of self-emulsifying system containing indomethacin to rats. Int J Pharm, 194(1)., 81-9.

24. L Ozguney, L Ozcan, G Ertan and T Guneri (2007), The Preparation and evaluation of sustained Release Suppositories of ketoprofen and Eudragit RL 100 by using Factorial design. Pharm Dev Tech, 12., 97-107.
25. Lintz W, Barth H, Osteloh G, Schmidt, Bothelt E (1998). Pharmacokinetics of tramadol and bioavailability of enteral tramadol formulation. 3rd Communication: Suppositories. Arzneimittelforschung, 48(9)., 889-99.
26. Malladi Shastry SP, Satyanarayana NV, Prakash V Diwan (1992), Krishna DR. Studies on the stability of fast and slow release compressed propranolol HCl suppositories. Indian. J Pharm Sci, 54(6)., 222-6.
27. Nair L and Bhargava HN (1999). Comparison of *in vitro* dissolution and permeation of fluconazole from different suppository bases. Drug Develop Ind Pharm, 25(5)., 691-4.
28. Nishimura K, Hidaka R, Hirayama F, Arima H, Uekama K (2006). Improvement of Dispersion and release properties of Nifedipine in suppositories by complexation with 2-hyroxypropyl-β-cyclodextrin, J.Incl.Phen and Macrocyclic Chem. 56(1-2)., 85-88.
29. Pandit JK, Choudhary PK and Mishra B (1990). *In vitro* evaluation of pentazocine suppositories. The Eastern Pharmacist, Jan., 171-3.
30. Ramarao P, Vijay Kumar D, Diwan PV(1998). Comparative pharmacokinetic evaluation of compressed rectal suppositories of diltiazem hydrochlorid. Ind J Pharmacol, 30(3)., 191-4.
31. Realdon N, Ragazzi E (2001). Effect of drug solubility on *in vitro* availability rate from suppositories with polyethylene glycol excipients. Pharmazie, 56(2)., 163-7.
32. Regdon G, Schirm S, Regdon G(1995). Formulation and *in vitro* study of antimalarial rectal suppositories. Pharmazie, 50., 439-440.
33. Regdon G, Schirm S, Regdon G(1996). Formulation and *in vitro* study of theophylline containing suppositories. Pharmazie, 51., 346-52.
34. Rishiraj C, Harsha B.Patel, Rajan DS, Jayanthi C, Suresh B(1999), Formulation and *in vitro* characterization of chloroquine phosphate suppositories. The Eastern Pharmacist, 4., 117-8.
35. Ryu JM, Chug SJ, Lee MH, Kim CK, Chang KS(1999). Increased bioavailability of propranolol in rats by retaining thermally gelling liquid suppositories. J Control Release, 59 (2)., 163-72.

36. S Maity, B.Sa, A k Bandyopadhay(2008). Development and evaluation of matrix and two layered sustained release suppositories of theophylline. Boll Chim Farm, 140(6)., 462-6.
37. Sastri MS, Satyanarayana MV, Krishna DR, Diwan PV(1993) *In vitro* and *in vivo* studies on slow release propranolol hydrochloride suppositories. Pharmacokinetic and pharmacodynamic evaluation. Arzneimittel-Forchung, 43(3)., 320-3.
38. Schmitt M and Guentert TW (1990). Influence of hydrophilicity of suppository bases on rectal absorption of carprofen, a lipophilic non-steroidal anti-inflammatory drug. J Pharm Sci, 79(4)., 359-63.
39. T A Adegboye, O Itiola (2008), Physical and Release Properties of Metronidazole Suppositories. Trop J. Pharm Res, 7(1) ., 887-896
40. Taneja LM, Singh J and Jaiswal SB. Bioavailability of insulin from insulin suppositories to rabbits. 1981; Vol. XXIV No. 288: 117-20
41. Tatsumi A, Oda S, Nakamoto T, Muraoka R, Takahashi V, Tanaka k et al (2008). Release control of prednisolone from suppository prepared using pulverized tablet. Yakugaku Zassi , 128(4)., 641-8.
42. Uzunkaya G, Bergladi N(2003). *In vitro* drug liberation and kinetics of sustained release indomethacin suppository. IL Farmaco, 58(7)., 509-12.
43. Yong CS, Yang CH, Rhee JD, Lee BJ, Kim DC, Kim DD et al(2004). Enhanced rectal bioavailability of ibuprofen in rats by poloxamer 188 and menthol. Int J Pharm, 269(1)., 169-76.